arXiv:1409.8385v1 [math.CA] 30 Sep 2014

Novel representation of the general Heun's functions. Back to the beginning

Plamen P Fiziev

Sofia University Foundation for Theoretical and Computational Physics and Astrophysics, Boulevard 5 James Bourchier, Sofia 1164, Bulgaria

and

BLTF, JINR, 141980 Dubna, Moscow region, Russia;

e-mail: fizev@phys.uni-sofia.bg and fizev@theor.jinr.ru

We study a novel type of solutions of the general Heun's equation, based on its symmetric form. We derive the symmetry group of this equation which is a proper extension of the Mobius group. The new series solution treat simultaneously and on an equal footing all singular points.

1 Introduction

The popular form of the general Heun's equation

$$H'' + \left(\frac{\gamma_{_G}}{z} + \frac{\delta_{_G}}{z-1} + \frac{\epsilon_{_G}}{z-a_{_G}}\right) H' + \frac{\alpha_{_G}\beta_{_G} z - \lambda}{z(z-1)(z-a_{_G})} H = 0, \quad \gamma_{_G}+\delta_{_G}+\epsilon_{_G} = \alpha_{_G}+\beta_{_G}+1;$$

treats asymmetrically its regular singular points $0, 1, a_{_G}, \infty$ with indices [1, 2, 3]

$$\{0, 1-\gamma_{_G}\}, \quad \{0, 1-\delta_{_G}\}, \quad \{0, \gamma_{_G}+\delta_{_G}-\alpha_{_G}-\beta_{_G}\}, \quad \{\alpha_{_G}, \beta_{_G}\}. \quad (1)$$

This causes difficulties in numerical calculations outside the circle $|z| < 1$, $z \in \mathbb{C}$, especially for $|z| \gg 1$.

The symmetric form of the general Fuchsian equation [4, 5] when applied to the special case of four regular singular points $z_{j=1...4} \in \mathbb{C}$ with the indices $\{\alpha_j, \beta_j\} : j = 0, ..., 4$, $\sum_{j=1}^{4}(\alpha_j + \beta_j) = 2$ is

$$\mathcal{W}'' + \left(\sum_{j=1}^{4} \frac{1-\alpha_j-\beta_j}{z-z_j}\right) \mathcal{W}' + \frac{1}{P(z)} \left(\lambda + \sum_{j=1}^{4} \frac{q_j}{z-z_j}\right) \mathcal{W} = 0, \quad (2)$$

$$P(z) = \prod_{j=1}^{4}(z-z_j) = \sum_{n=0}^{4}(-1)^n \sigma_{_{4-n}} z^n, \qquad q_j = \alpha_j\beta_j P'(z_j) \ \text{ for } \ j = 1...4. \quad (3)$$

Making use of the transformation

$$\mathcal{W}(z) = \mathcal{F}(z) \prod_{j=1}^{4} (z-z_j)^{\nu_j}, \ \sum_{j=1}^{4} \nu_j = 0, \quad (4)$$

and imposing *symmetric* constraints on the indices

$$\alpha_j + \beta_j = \tfrac{1}{2}, \ \alpha_j\beta_j P'(z_j) = q_j, \ \text{for} \quad j = 1..4, \quad (5)$$

we obtain the following simple symmetric form of the general Heun's equation

$$\mathcal{F}'' + \frac{1}{2}\left(\sum_{j=1}^{4} \frac{1}{z-z_j}\right) \mathcal{F}' + \frac{1}{P(z)} \left(\lambda + \sum_{j=1}^{4} \frac{q_j}{z-z_j}\right) \mathcal{F} = 0, \quad (6)$$

or in a self-adjoint form:

$$(P(z))^{1/2} \left((P(z))^{1/2} \mathcal{F}'\right)' + (\lambda + Q(z)) \mathcal{F} = 0, \quad (7)$$

where

$$Q(z) = \sum_{j=1}^{4} \frac{q_j}{z-z_j} = -\tfrac{1}{16P(z)} \sum_{j=1}^{4} (\sin(2\chi_j))^2 \, \partial_z P(z_j) \partial_{z_j} P(z) \quad (8)$$

Here we use new uniformization parameters $\chi_{j=1..4} \in \mathbb{C}$:

$$\alpha_j = \tfrac{1}{2}(\cos\chi_j)^2, \quad \beta_j = \tfrac{1}{2}(\sin\chi_j)^2,$$
$$q_j = \tfrac{1}{16}(\sin(2\chi_j))^2 \prod_{k\neq i}^{4} (z_j - z_k).$$

2 The orthogonality and the invariance group of solutions

Proposition 1: For any two solutions $\mathcal{F}_{\lambda_{1,2}}(z)$ of the Eq. (7) with $\lambda_1 \neq \lambda_2$ we have

$$\int_{\mathcal{L}_{ij}} \mathcal{F}_{\lambda_1}(z)\mathcal{F}_{\lambda_2}(z)d\mu(z) = 0,$$
$$d\mu(z) = (P(z))^{-1/2}\, dz. \qquad (9)$$

Here $\mathcal{L}_{ij} \in \mathbb{C}$ is any contour which starts at the singular point z_j and ends at the singular point z_j without going through the other singular points $z_{k\neq i,j}$. Besides, the singular boundary conditions

$$(P(z))^{1/2}\left(\mathcal{F}_{\lambda_1}(z)\mathcal{F}'_{\lambda_2}(z) - \mathcal{F}_{\lambda_2}(z)\mathcal{F}'_{\lambda_1}(z)\right) \restriction_{z_j,z_j} = 0 \quad (10)$$

are supposed to be satisfied. ◀

As seen, in our symmetric formulation of the general Heun's equation (7) the parameter λ plays natural role of eigenvalue in the corresponding two-singular-point boundary problems for a self-adjoined operator [2, 3, 8, 9]. This is also consistent with the profound results of [10].

Proposition 2: Equation (**??**) is invariant under the *extension* $\widehat{\mathfrak{G}}_{Mobius}$ of the standard Mobius group $\mathfrak{G}_{Mobius}$. This extension acts on the functions of 10 variables $\mathcal{F}(z; z_1, ..., z_4; q_1, ..., q_4; \lambda)$ and is produced by the following basic transformations:

(i) Complex translations with $\zeta \in \mathbb{C}$:

$$z \to z + \zeta; \quad z_j \to z_j + \zeta : j = 1, ..., 4;$$
$$q_j \to q_j : j = 1, ..., 4; \quad \lambda \to \lambda. \qquad (11)$$

(ii) Complex dilatations with $t \in \mathbb{C}$, $t \neq 0$:

$$z \to t\, z; \quad z_j \to t\, z_j : j = 1, ..., 4;$$
$$q_j \to t^3 q_j : j = 1, ..., 4; \quad \lambda \to t^2\lambda. \qquad (12)$$

(iii) Inversion

$$z \to 1/z; \quad z_j \to 1/z_j : j = 1, ..., 4;$$
$$q_j \to -q_j/\left(z_j^2\sigma_4\right) : j = 1, ..., 4;$$
$$\lambda \to \left(\lambda - \sum_{j=1}^{4} q_j/z_j\right)/\sigma_4. \qquad \blacktriangleleft \quad (13)$$

Using proper compositions of these basic transformations (see [10, 11] for $\mathfrak{G}_{Mobius}$) we are able to construct a representation of the whole extended Mobius group $\widehat{\mathfrak{G}}_{Mobius}$ that acts on the solutions $\mathcal{F}(z; z_1, ..., z_4; q_1, ..., q_4; \lambda)$ of Eq. (**??**) without bringing us outside of the variety of these solutions. Hence, the extended group $\widehat{\mathfrak{G}}_{Mobius}$ is the group of invariance of the variety of solutions to Eq. (**??**).

3 The Taylor series expansion of the solutions

Our next step is to adopt the following basic assumption which is of crucial importance for further work:

$$z_{j=0,...,4} \neq 0. \qquad (14)$$

Then the function $\mathcal{F}(z)$ is an analytical one in the vicinity of the point $z = 0$ and has an absolutely convergent Taylor series expansion

$$\mathcal{F}(z) \equiv \mathcal{F}(z; z_1, ...z_4; q_1, ...q_4; \lambda) = \sum_{n=0}^{\infty} f_n(z_1, ...z_4; q_1, ...q_4; \lambda)z^n \quad (15)$$

with the coefficients $f_n(z_1, ...z_4; q_1, ...q_4; \lambda)$ defined by the nine-term recurrence relation

$$f_n + \sum_{k=1}^{8} r_{n-k} f_{n-k} = 0, \qquad (16)$$

which can be obtained from Eqs. (**??**) and (**??**).

We shall take advantage of the freedom to put the singular points $z_{j=0,...,4}$ in the proper places in the complex plane $\mathbb{C}$ for simplifying, as much as possible, the coefficients in recurrence (16) and thus, the very solutions $\mathcal{F}(z)$. This can be done in a symmetric way by imposing additional conditions on the elementary symmetric functions $\sigma_{j=1,2,3,4}$. Using the proper Mobius transformation one is able to impose three independent constraints on $z_{j=0,...,4}$ without changing the problem. Obvious simple choice is to reduce the quartic equation $P(z) = 0$ to the following biquadratic one: $z^4 - 2\cos(2\phi)z^2 + 1 = 0$ with the roots

$$z_1 = e^{i\phi},\ z_2 = -e^{-i\phi},\ z_3 = -e^{i\phi},\ z_4 = e^{-i\phi}, \quad (17)$$

by imposing three symmetric constraints

$$\sigma_1 = \sigma_3 = 0,\ \sigma_4 = 1, \qquad (18)$$

and replacing $\sigma_2 = -2\cos(2\phi)$ with one more complex uniformization parameter $\phi \in \mathbb{C}$.

The meaning of the new variable ϕ is revealed by the formula for the invariant $a(z_1, z_2, z_3, z_4)$ of the Mobius transformation – the so called *cross-ratio*. In our problem it acquires the form

$$a = \frac{(z_1 - z_3)(z_2 - z_4)}{(z_2 - z_3)(z_1 - z_4)} = \frac{1}{(\sin\phi)^2} \quad \Rightarrow$$
$$\sigma_2 = -2\left(1 - \frac{2}{a}\right). \quad (19)$$

As a result, one obtains for the coefficients in recurrence (16) in the following form:

$$r_{n-1} = 0, \quad (20)$$
$$r_{n-2} = \tfrac{1}{n(n-1)}\Big(\lambda - \tfrac{i}{4}\,\sin(2\phi)\,\rho_2\Big) + 4\left(1 - \tfrac{5}{n} + \tfrac{3}{2}\tfrac{1}{n-1}\right)\cos(2\phi), \quad (21)$$
$$r_{n-3} = -\tfrac{1}{n(n-1)}\tfrac{i}{4}\,\sin(2\phi)\,\rho_3, \quad (22)$$
$$r_{n-4} = \tfrac{1}{n(n-1)}\Big(-2\lambda\cos(2\phi) + \tfrac{i}{4}\,\sin(2\phi)\,\rho_4\Big) + 2\left(1 - \tfrac{16}{n} + \tfrac{9}{n-1}\right)\left((\cos(2\phi)^2 + 1\right), \quad (23)$$
$$r_{n-5} = \tfrac{1}{n(n-1)}\tfrac{i}{4}\,\sin(2\phi)\,\rho_5, \quad (24)$$
$$r_{n-6} = \tfrac{\lambda}{n(n-1)} - 4\left(1 - \tfrac{33}{n} + \tfrac{45}{2}\tfrac{1}{n-1}\right)\cos(2\phi), \quad (25)$$
$$r_{n-7} = 0, \quad (26)$$
$$r_{n-8} = 1 - \tfrac{56}{n} + \tfrac{42}{n-1}. \quad (27)$$

where we introduce the following four functions of five variables $\{\phi, \chi_1, \chi_2\chi_3, \chi_4\}$:

$$\rho_2 = \Big((\sin(2\chi_1))^2 + (\sin(2\chi_3))^2\Big) - \Big((\sin(2\chi_2))^2 + (\sin(2\chi_4))^2\Big),$$
$$\rho_3 = e^{-i\phi}\Big((\sin(2\chi_1))^2 - (\sin(2\chi_3))^2\Big) + e^{i\phi}\Big((\sin(2\chi_2))^2 - (\sin(2\chi_4))^2\Big),$$
$$\rho_4 = e^{2i\phi}\Big((\sin(2\chi_1))^2 + (\sin(2\chi_3))^2\Big) - e^{-2i\phi}\Big((\sin(2\chi_2))^2 + (\sin(2\chi_4))^2\Big),$$
$$\rho_5 = e^{i\phi}\Big((\sin(2\chi_1))^2 - (\sin(2\chi_3))^2\Big) + e^{-i\phi}\Big((\sin(2\chi_2))^2 - (\sin(2\chi_4))^2\Big).$$

4 The circular case

There exist two quite different cases of positions of the singular points $z_{j=0,\dots,4}$.

The first (general) case is the one in which the singular points $z_{j=0,\dots,4}$ do not lie on any circle in the complex plane. We call it *the non circular case*. In the non circular case, the theory of solutions of Eq. (**??**) is more complicated. It is not enough developed for the popular form of the general Heun's equation with complex $a_{_G} \notin \mathbb{R}$, see the Introduction.

The second (special) case is the one when all regular singular points $z_{j=0,\dots,4}$ of Eq. (**??**) lie on some circle $\mathfrak{C} \in \mathbb{C}$. We will call this case *the circular case*. A particular and natural such case is the one when $z_{j=0,\dots,4} \in \mathbb{R}$, i.e., all singular points are real.

In the circular case, one is able to move all singular points $z_{j=0,\dots,4}$ on any other circle $\tilde{\mathfrak{C}} \in \mathbb{C}$ using Mobius transformation which preserves the circular property and transforms any circle in the complex plane into another circle, see [10, 11]. Especially, in [10] the circular case for the general Heun's equation with $a_{_G} \in \mathbb{R}$ was substantially elaborated using the standard choice (**??**) and without any relation with the choice (17) in Eq. (**??**).

The value of the invariant cross-ratio (**??**) $a \in \mathbb{R}$ is real for any four complex points $z_{j=1,2,3,4}$ on a circle in $\mathbb{C}$. Then, from relation (**??**) follows that in the circular case the angle $\phi \in \mathbb{R}$ is real and the singular points (17) lie on the unit circle with the center $z = 0$.

Thus, according to the basic results of the standard analytic theory of ordinary differential equations [6, 7], we obtain our key result:

Proposition 3: In the circular case, the series (**??**) with coefficients (**??**)-(**??**) and $\phi \in \mathbb{R}$ are absolutely convergent inside the unit circle, i.e., for any $z \in \mathbb{C}$ with $|z| < 1$. ◀

Corollary: In the circular case, the four regular singular points $z_{j=1,2,3,4}$ of the general Heun's equation (**??**) can be treated simultaneously and equally from inside the unit circle using the Taylor series (**??**).

Next important step is to restrict Proposition 4 (iii) to the circular case.

Proposition 4: In the circular case, equation (**??**) preserves its form if one makes the following substitutions

$$\begin{aligned} z &\to 1/z, \quad z_j \to 1/z_j, \\ F(z) &\to F(1/z), \quad P(z) \to P(1/z), \\ \lambda &\to \lambda - \tfrac{i}{4}\sin(2\phi)\rho_2, \quad q_j \to -q_j/z_j^2. \end{aligned} \quad (28)$$

This way we obtain from solutions (**??**) new solutions of Eq. (**??**) in the form of the Laurent series

expansions which are absolutely convergent for any $z \in \mathbb{C}$ with $|z| > 1$. ◀

Corollary: In the circular case, the four regular singular points $z_{j=1,2,3,4}$ of the general Heun's equation (**??**) are mapped under inversion on the same points, removed to the initial positions $z_{j=4,3,2,1}$, respectively. Hence, one can treat all singular points simultaneously and equally from outside the unit circle using the corresponding Laurent series, described in Proposition 6.

As a final result, in the circular case we reach a totally symmetric treatment of the singular points $z_{j=1,2,3,4}$ in the whole complex plane $\tilde{\mathbb{C}}$.

5 Concluding remarks

In the present talk, we introduced and studied a novel representation of the general Heun's functions. It is based on the symmetric form of the Heun's differential equation yielded by a further development of the Felix Klein symmetric form of the Fuchsian equations with an arbitrary number $N \geq 4$ of regular singular points.

The basic relations for the general Heun's equation with $N = 4$ are derived and discussed in detail.

We derived the symmetry group of this equation and its solutions. It turns to be a proper extension of the Mobius group.

Special attention was paid to the nine-term recurrence relation for the coefficients of the Taylor series solutions of the novel symmetric form of the general Heun's equation.

We also showed that the novel Taylor series solutions are absolutely convergent inside the circle with the unit radius. After the simple inversion of the independent variable $z \to 1/z$ one obtains also the Laurent series solutions which are absolutely convergent outside the circle with unit radius.

Since the four regular singular points of the symmetric form of the general Heun's equation lie on the unit circle, one can use the new solutions for a simultaneous equal treatment of all of them.

One can hope that this new approach will simplify the solution of the existing basic open problems in the theory of the general Heun's functions.

The novel representation will allow also development of new effective computational methods for calculations with the Heun's functions which at present are still a quite problematic issue.

Analogous results for the four cases of the confluent Neun's equation will be discussed elsewhere.

Acknowledgements

The author is grateful to Professor Sergei Yu. Slavyanov for his helpful comments, help with the literature, and his kind encouragement, as well as to Professors Alexander Kazakov and Artur Ishkhanyan for their interest in the basic results of the present talk. Special thanks to Proffesor Oleg Motygin for careful reading of manuscript and useful remarks.

Special thanks are also to the leader of the Maple-Soft-developers, Edgardo Cheb-Terrap, for many useful discussions during the last years on the properties of the Maple-Heun's functions and the computational problems with them.

The author is also thankful to the leadership of BLTP, JINR, Dubna for the support and good working conditions. This talk was also supported by the Sofia University Foundation *Theoretical and Computational Physics and Astrophysics* and by the Bulgarian Agency for Nuclear Regulation, the 2014 grant.

References

[1] Heun K 1889 *Math. Ann.* **33** 161

[2] Ronveaux A (ed.), 1995 Heun's Differential Equations, Oxford Univ. Press, New York

[3] Slavyanov S Y, Lay W 2000 Special Functions, A Unified Theory Based on Singularities (Oxford: Oxford Mathematical Monographs)

[4] Klein F 1894 Forlesungen über lineare Differentialgleihungen der zweiten Ordnung (Göttingen)

[5] Forsyth A R 1902 Theory of Differential Equations Part III (Cambridge University Press)

[6] Golubev V V 1950 Lectures on the analytic theory of differential equations, Moscow (in Russian)

[7] Coddington E A, Levinson N 1955 Theory of Ordinary Differential Equations (TMH Eddition 1972, 9th Reprint 1987, Mc Graw-Hill Inc. N Y)

[8] Kazakov A Ya , Lay W, Slavyanov S Yu 1996 Eigenvalue prolems for Heun's equations, *Algebra and Analysis*, **8** 129-141 (in Russian)

[9] Lay W, Slavyanov S Yu 1998 The central two-point connection problem for the Heun class of ODE *J Phys A: Math Gen* **31** 8521-8531

[10] Smirnov V I (1996) Selectet papers. Analytical theory of ordinary differential equations (St. Petersburg University Press) (in Russian)

[11] Ablowitz M, Fokas A S 2003 Complex Variables Introduction and Applications (Cambridge University Press)